

Horticultural News

***Volume 100, Number 4
Fall, 2020***



Horticultural News

Editors: Winfred P. Cowgill, Jr. & Wesley R. Autio

The New Jersey State Horticultural Society was organized on August 17, 1875 at Geological Hall, Rutgers College, New Brunswick, NJ. It remains the oldest Horticultural organization in New Jersey.

Horticultural News began as the *The New Jersey State Horticultural Society News*, in October of 1920. The Society began "collecting paid membership in order to obtain funds to promote new features of the society and extend the usefulness of the society. The Horticultural Society News was started to be the official society publication." Published M. A. Blake, Professor at Rutgers College was the first president and chair of the publication committee.



Editors served as follows:

MA Blake	1920 - 1947
Norman F Childers	1948 - 1980
Win Cowgill	1981 - 1988
Emily Brown Rosen	1988 - 1990
Linda Butenis Vorsa	1991 - 1995
Jerry Frecon	1995 - 2010

June 2010: *Horticultural News* has moved to an online web-based format. The New Jersey State Horticultural Society has partnered with the University of Massachusetts *Fruit Notes*, Dr. Wesley Autio, Editor. Cowgill and Autio will be the new editors of *Horticultural News* and *Fruit Notes*.

Horticultural News is distributed to growers, extension personnel and researchers and libraries across North America. Horticultural News focuses primarily on tree-fruit culture, but addresses small-fruit cultural issues as well. Most reports are from current research at Rutgers University, University of Massachusetts, and other universities.

Horticultural News is published four times per year by the New Jersey State Horticultural Society. It is provided as a benefit to membership in the society. Membership costs \$50 per year. Each one-year subscription begins January 1 and ends December 31. Payments via check must be in United States currency and should be payable to the New Jersey State Horticultural Society. Horticultural News Electronic subscriptions are available as benefit of membership in the NJ State Horticultural Society. A hidden link will be mailed immediately after publication to members. Issues will be made freely available on this website six months after publication.

Correspondence should be sent to: New Jersey State Horticultural Society

Greg Donaldson
176 Airport Road
Hackettstown, NJ 07840
TEL (908) 296-1064

For advertising opportunities, contact: Win Cowgill: wincowgill@mac.com; (908) 489-0207

Table of Contents

Massachusetts Fruit IPM Report, 2020

Jaime Piñero, Daniel Cooley, Jon Clements, Sonia Schloemann, and Elizabeth Garofalo 1

Early-season Performance of Diluted Concord Grape Juice and Commercial Lures at Attracting Spotted Wing Drosophila and Effects of Juice Fermentation on Trap Captures

Jaime C. Piñero and Heriberto Godoy-Hernandez..... 10

Thwarting the Low-spray Plan: Marssonina Throws a Curve Ball

Elizabeth Garofalo and Daniel Cooley 16

Northern New Jersey Report, As Presented at the NEIPM Conference in October 2020

Win Cowgill..... 21

New Jersey News 24

Cover: UMass Cold Spring Orchard Research & Education Center Photo: Wes Autio.

Massachusetts Fruit IPM Report, 2020


Jaime Piñero, Daniel Cooley, Jon Clements, Sonia Schloemann, and Elizabeth Garofalo
University of Massachusetts

Weather

Other than some late fall 2019 cold and an early December snowstorm, the winter of 2019-20 was characterized by mild temperatures and lack of snow. The minimum (at the UMass Orchard, Belchertown, MA) was 2.4 degrees F. on 15-February, 2020. Given the lack of snow cover and some early March warmth, spring 2020 seemed on the brink of being “early,” but April turned cold after early green tip arrived on 1-April. Mid-April temperatures flirted with bud damaging lows, and in fact it is suspected some site-specific bud injury occurred during this time. Full bloom was about 16-May, and it was extended with cool May weather. May and June were fairly dry which was great for controlling scab but initiated drought conditions in MA. July was hot, very hot overall, with a maximum temperature of 94.2 on 19-July.

A quick analysis of 20 years of weather station data from the UMass Orchard makes it pretty clear July has been getting warmer over the period from 2001-2020. (Fig. 1) At the UMass Orchard July and August had 3.90 and 4.77 inches of rain respectively during thunderstorms, which ameliorated our drought situation somewhat. However, many other orchards in MA were extremely dry and without irrigation fruit size suffered, and with irrigation, growers and water supplies were run ragged. Hailstorms were out there this summer, and several orchards had extensive hail damage. Crop insurance is an essential risk management strategy these days. Finally, the weather was generally sunny late summer into early fall, and peach and apple quality was very good as a result.

The fall harvest season in particular was weather-benevolent, and customer turnout at PYO orchards and farm stands could not have been much better on account






Figure 2. (A) Characteristic fireblight ooze on an infected apple shoot. This infection originated with a blossom infection resulting from late bloom time and. Streptomycin was applied to earlier blooming varieties but this late bloomer was missed, (B) advanced shoot blight with darker ooze droplets, (C) later in the season, fireblight persisted in the cider block and affected developing fruit as well as shoots.

of Covid-weary people wanting to get outside and enjoy a safe experience with their family. But growers had to adapt by adopting significant Covid-19 safety measures (at their expense) such as a customer reservation system, touchless payment, social distancing signage, requiring face masks, and hiring extra staff to manage the situation. Most growers will be extra glad to see the 2020 growing season come to a close, as whereas weather used to be the big worry item, they had an additional worry item with Covid-19 in 2020.

Diseases

Apple scab was largely a no-show as a result of the drought that eventually expanded to engulf the entire state. Final ascospores were observed in the petri plate assay and funnel trap on June 29, 2020. The cool temperatures heading into bloom led us to think we might slip through 2020 with no **fire blight**, but, sudden increases in temps throughout the region led to outbreaks in late blooming varieties. Suspect fire blight samples from five orchards throughout Massachusetts were submitted to Anna Wallis, PhD. student in Kerik Cox lab at Cornell University, for confirmation and streptomycin resistance testing. All were positive for fireblight and negative for resistance to streptomycin.

Sooty blotch (SB) and **flyspeck (FS)** became apparent in the harvest surveys in several blocks. For example, in two blocks located in a single MA orchard having cider apple cultivars, the incidence of SB and

FS were 3.7 and 9.6%, respectively for block one, and 0% (SB) and 1.8% (FS) for the second block.

Marssonina leaf blotch continues to creep up in many orchards, cultivar and management practices significantly influence the severity of the disease. However, even in rigorously managed conventional orchards, the disease has been observed. The following photos (Figs. 3 and 4) all were taken in a low spray



Figure 3. Early *Marssonina* symptoms. On close inspection acervuli (tiny black fruiting fungal bodies) can be seen in the brown-purple spots. Photo taken in Amherst, MA October 9, 2020.



Figure 4. Enterprise (front of photo) versus GoldRush (behind) leaf drop.

organic orchard block (no fungicides this year).

Insects

In 2020, the two most damaging insect pests were **plum curculio (PC)** and **tarnished plant bug (TPB)**. Across 11 commercial orchards located in MA (7), NH (3), and ME (1), 10,560 fruits were inspected for insect injury at harvest. The average level of block-wide injury was 2.8% for PC and 2.6% for TPB. No PC research involving odor-baited trees was conducted this year. Therefore, the incidence of injury by PC being reported here occurred under standard grower manage-

ment. As shown in Table 1 below, injury by internal Lepidoptera was very low. In 2019, low levels of fruit injury attributable to codling moth (CM), Oriental fruit moth (OFM) and obliquebanded leafrollers (OBLR) in Massachusetts were recorded: 0%, 0.15%, and 0.08%, respectively.

Plum curculio (PC). The timing of the petal fall insecticide spray is critical. There were at least 3 orchards where the petal fall spray was applied later than growers intended. This was likely due to the presence of mixed cultivars in those blocks or to rainy weather conditions that may have prevented them from entering the blocks to spray. In one block located in the UMass Cold Spring

Table 1. Insect incidence at harvest in 10,560 apples, sampled from 11 orchards located in MA (7), NH (3), and ME (1).

	PC	Stink bug	TPB	AMF	PC feeding or other damage	Rollers	OFM	CM	EAS	San Jose scale
PERIMETER	4.28	0.17	1.76	0.31	0.11	0.09	0.06	0.20	0.01	0.11
INTERIOR	1.27	0.45	3.52	0.54	0.11	0.06	0.03	0.26	0.31	0.00
Whole orchard block	2.78	0.31	2.64	0.43	0.11	0.07	0.04	0.23	0.16	0.06

Orchard that was used to compare the level of control achieved with Verdepryn (a.i. Cycloprothrin) versus that of Avaunt (a.i. Indoxacarb), the insecticides were sprayed a couple of days after the optimal time. While both insecticides were equally effective at controlling PC, the timing of application led to higher levels of PC injury in those blocks, compared to other orchard blocks.

Monitoring for PC is key to successful management. A monitoring technique, based solely on observation of fresh PC injury on fruit from odor-baited trees, has proven effective and efficient at determining the need for and appropriate timing of perimeter-row insecticide sprays against PC after the whole-block petal fall spray. A [Fact Sheet](#) is available here.

Tarnished plant bug (TPB). Injury by this insect pest was recorded at each orchard where we conducted harvest surveys. This year, we initiated a project (led by UMass graduate student Ms. Prabina Regmi) aimed at evaluating plant volatiles for potential use in monitoring systems for TPB. *Findings will be presented in the next Fruit Notes issue.* Research will continue in 2021.

Apple maggot fly (AMF). In 2020, we conducted a study that started in 2019 in collaboration with the Univ. of New Hampshire and the Univ. of Maine. This investigation, led by a UMass graduate student (Ms. Dorna Saadat) evaluated the efficiency of an attract-and-kill strategy involving the use of attractive lures and insecticide sprays in combination with sugar added as phagostimulants, applied to perimeter-row trees. Two treatments were compared: (1) attract-and-kill and (2) grower control. The attract-and-kill block involved the use of lures deployed on the perimeter of the block plus red spheres as a monitoring tool. In these blocks, perimeter-row trees were sprayed with insecticide mixed with 3% sugar added to the tank mix. In the grower control blocks, two or three insecticides were applied to the entire block by the grower. *A full account of the results will be published in the next issue of Fruit Notes.*

Internal Lepidoptera. In 2020, we conducted a study aimed at assessing the efficacy of mating disruption using dual pheromone lure dispensers targeting obliquebanded

leafroller (OBLR) and codling moth (CM). This work was led by a UMass graduate student, Mr. Ajay Giri. The mating disruption dispensers were deployed at a low rate (32 dispensers/acre). At harvest, we quantified the level of injury by CM and OBLR in the mating disruption and in the grower control blocks. The average level of fruit injury by CM recorded in the three mating disruption (MD) and the three grower control (GC) blocks was 0.02% and 0.03% respectively. Injury by OBLR was similarly low: 0.08% and 0.15% in MD and GC blocks, respectively. *More detailed results will be published in the next issue of Fruit Notes.*

Mites. Mites were not reported by growers as being a problem despite the lack of rain.

Brown Marmorated Stink Bug. Monitoring of BMSB was done in 10 commercial orchards starting in mid-August, 2020. Six of those locations had not been monitored for BMSB before. BMSB was detected at every single orchard that was monitored, indicating the presence of established populations state-wide. Ghost traps (insecticide-treated netting deployed in association with the BMSB pheromone lure) were deployed at selected orchards. In one orchard, two ghost traps killed 636 BMSB in 3 weeks. We can conclude that BMSB populations continue to be on the rise, and injury by this invasive pest was reported in at least 3 Massachusetts orchards.

San Jose scale was detected in only a couple of orchards, and the level of injury was low.




Figure 5. Injury by apple leaf midge and midge adults (insert).




Figure 6. Carbohydrate balance at UMass Orchard culminating in a deficit of nearly 100 g/day on 26-May.

Injury by **European apple sawfly (EAS)** was recorded in a couple of orchards, at low levels.

Apple leaf curl midge. The apple leaf midge (also known as apple leaf curling midge and apple leaf gall midge) is an exotic foliar pest of apple trees. It was found in a few MA orchards. Minor injury potentially caused by this insect pest was suspected in one orchard.

Horticulture

Two things stand out in 2020. First, the weather, or should I say the carbohydrate deficit during the 10 mm apple fruitlet chemical thinning stage, and second, wind during Tropical Storm Isaias which wreaked havoc on some apple trees at the UMass Orchard.

2020 Thinning weather — suffice it to say a rather massive carbohydrate deficit on account of warm day and night temperatures and cloud cover made chemical thinning in 2020 rather dubious. More and more we encourage growers to thin aggressively using a “nibble” approach of multiple chemical thinning applications beginning at bloom. But still, the single best time to accomplish chemical thinning is when apple fruitlets are about 10 mm. This year, a large carbohydrate deficit occurred about this time, culminating in a 4-day average carbohydrate balance of -95 on 26-May. (Fig. 6) This is a huge deficit and could easily result in over-thinning if a chemical thinner was applied. The deficit was short-lived, but there was much hollering by growers who did apply a chemical thinner just prior to this deficit. After the fact a resounding “oh my goodness, I think I thinned all my fruit off!” was heard. We have not experienced this situation in several years. Some varieties

did turn out light, however, later blooming apples had a nice crop. Mac type apples were generally good, but Honeycrisp turned out rather light across the board. Lesson learned? Pay attention to the carbohydrate deficit during that time period when apple fruitlets are about 10 mm, adjust your chemical thinning response (time, rate, combinations) accordingly, and direct 2/3 of the thinning spray in the tops of trees (always).

Wind and G.41 are a mismatch — Tropical storm Isaias blasted through the Hudson Valley of NY on 04-August with heavy rain there, but ALL we got at the UMass Orchard in Belchertown was WIND, with a recorded on-site gust of 50 mph. ALL was enough to break off many (100 or so, I did not count exactly, it was too depressing) young apple trees at the Orchard. The consistent factor was all the broken apple trees were on G.41 rootstock, they were snapped right off at the graft union. (Fig. 7) It appeared to me that this was a result of several factors: moderate-heavy crop loads, long branches, and a marginal support system with failure to fully keep the trees from twisting or moving too much from vertical along with what is commonly known as a weak graft union when mated with G.41 rootstock. Some varieties, but not all are known, may exacerbate the problem. Young trees are also more prone to breakage. Lesson learned? Apple trees planted on G.41 rootstock cannot be allowed to twist or rock in the wind, period. Keep branches short. Build a proper support system. Use extra fasteners (or wires) when attaching trees to wire. G.41 is a good rootstock because it is fire blight resistant, but it has this one fatal flaw that needs extra attention if planting an apple orchard on this rootstock. I have a YouTube video here if you



Figure 7. Apple tree on G.41 rootstock snapped right off at the graft union in high wind.

want to see the carnage up-front and personal: <https://youtu.be/XLXvrBFY6H4>.

Small Fruit IPM

Winter Moth (WM). Based on the success of biocontrols of Winter Moth in recent years and the resulting lower population levels, we did not directly monitor WM in 2020. Information on WM emergence predictions based on observations in RI and on Degree Day models were disseminated in the IPM Berry Blast and Healthy Fruit Newsletters early in the season.

Gypsy Moth (GM). Based on low levels of Gypsy Moth egg masses in the winter of 2019/2020, the UMass Fruit Extension team did not do any direct systematic monitoring of GM populations during the

2020 growing season. Information on damage potential in hot spots was disseminated in IPM Berry Blast and Healthy Fruit Newsletters as needed.

Spotted Wing Drosophila. In 2020, we continued to emphasize to implement the 1-2-3 IPM approach to spotted wing Drosophila (SWD) management. The three components are (1) monitoring, (2) cultural control methods, and (3) insecticide sprays. In 2020, we compared the performance of diluted Concord grape juice (1:3 juice: water) at capturing SWD against that of four commercial lures, at four MA locations. Diluted grape juice was used to monitor SWD for the entire season. The season-long pattern of SWD captured by diluted grape juice is presented in Fig. 8, below. *The results of this study are presented in the next article of this Fruit Notes issue.*

Massachusetts IPM Berry Blasts/Healthy Fruit Newsletter Small Fruit section. Nineteen issues of *Massachusetts IPM Berry Blast* (515 subscribers), were sent out during the 2020 growing season. These covered IPM recommendations for a wide range of pests and disease problems in berry crops. A con-




Figure 8. Season-long captures of adult spotted wing drosophila in traps baited with diluted concord grape juice (n= 4 orchards).

densed version of this information was also included in 20 issues of the Healthy Fruit newsletter (165 subscribers).

Special Projects

NEW Fruit IPM Fact Sheets and Videos! We have revamped the **Fruit IPM Fact Sheet series** (apple, blueberry, strawberry). If you are reading a PDF copy of this report, then the Fact Sheets are available (<https://ag.umass.edu/fruit/fact-sheets>). Otherwise, you can Google “UMass Fruit IPM Fact Sheets”. Educational **videos** relating to fruit production and pest management are available (https://www.youtube.com/channel/UUCKCU0_6fvuSPLtWvsmDhfwg/videos?view=0&sort=p&flow=grid) or Google “UMass Extension fruit program YouTube channel”.

Northeast Cider Apple Project (NECAP) -- Beginning in Fall 2019, this 3-year Project funded by NESARE is led by University of Vermont with collaborators from UMass and UMaine. At UMass Cooley, Pinero, Clements, and Garofalo are evaluating at least five cider orchards throughout Massachusetts for insect and disease incidence on cider apples. We are also evaluating horticultural and fruit quality characteristics to develop fact sheets and recommendations for both established and new growers of cider apples (<https://www.youtube.com/channel/UCWrmWfBqbcK8FgjVTuRT0Gw>).

MyIPM app -- work continued by Cooley, Clements, and Garofalo on the MyIPM including adding pear insects, cherry insects, and updating apple and pear diseases. MyIPM is designed to provide mobile access to pest management information for many fruit crops with an emphasis on resistance management. For more information on the app: <https://apps.bugwood.org/apps/myipmseries/>

Clements, J., D. Cooley, and E. Garofalo. A comparison of four on-site weather stations and one virtual weather service as data sources in 2020 for the apple scab infection period model at the UMass Cold Spring Orchard in Belchertown, MA. (Research/demonstration).

Clements, J., D. Cooley, and P. O’Connor. A comparison of using the ‘Ferri’ version of the fruitlet growth rate model and the Malusim app to predict fruit set in 2020 of Gala, Honeycrisp, Empire and Pazazz apples at the UMass Cold Spring Orchard in Belchertown, MA.

(Research/demonstration).

Clements, J. and J. Pinero. Blending technology and IPM: Onset Hobo RX300 weather station and NEWA, DTN Smart Traps, and “attract and kill” trap of brown marmorated stink bugs. A case study in a Berkshire’s Massachusetts orchard. (Research/demonstration).

Clements, J., D. Cooley, P. O’Connor, and L. Ware. OrchardWatch: remote sensing of weather conditions across multiple locations in a single orchard, does it make a difference? (Research/demonstration).

Publications

Clements, J. 2019. IFTA Tours Ontario. Fruit Notes 84:4 Fall 2019.

Piñero, J., D. Cooley, J. Clements, S. Schloemann, and E. Garofalo. 2020. Massachusetts Fruit IPM Report for 2019. Fruit Notes 85:1 Winter 2020.

Clements, J. 2020. Malusim App and Precision Apple Thinning – Trials and Tribulations. Fruit Notes 85:1 Winter 2020.

Clements, J. 2020. DTN Smart Traps – Worth it or Not? Fruit Notes 85:1 Winter 2020.

Clements, J. and E. Garofalo. 2020. Increasing Branching of Cider Apple Trees. Fruit Notes 85:1 Winter 2020.

Bradshaw, T., J. Clements, D. Cooley, E. Garofalo, J. Pinero, and R. Moran. 2020. New England Cider Apple Project. Fruit Notes 85:1 Winter 2020.

Clements, J., E. Garofalo, and W. Autio. 2020. 2015 Modi Organic NC-140 Apple Rootstock Trial and Drapenet Demonstration. Fruit Notes 85:3 Summer 2020.

Clements, J. J. Krupa, P. O’Connor, L. Ware, and D. Cooley. 2020. ‘OrchardWatch’ Weather Monitoring Grid at UMass Cold Spring Orchard. Fruit Notes 85:3 Summer 2020.

Usman, M., Gulzar, S., Wakil, W., Wu, S., Piñero, J.C., Leskey, T.C., Nixon, L.J., Oliveira-Hofman, C., Toews, M.D., and Shapiro-Ilan, D. 2020. Virulence of entomopathogenic fungi to the apple maggot *Rhagoletis*

pomonella (Diptera: Tephritidae) and interactions with entomopathogenic nematodes. Journal of Economic Entomology (in press).

Usman, M., Gulzar, S., Wakil, W., Piñero, J.C., Leskey, T.C., Nixon, L.J., Oliveira-Hofman, C., Wu, S. and Shapiro-Ilan, D. 2020. Potential of entomopathogenic nematodes against the pupal stage of the apple maggot *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae). Journal of Nematology e2020-79 | Vol. 52.

Piñero, J.C. and Manandhar, R. 2020. Ant attendance and arthropod diversity on elderberry extrafloral nectaries are influenced by plant genotype and pruning method. Arthropod-Plant Interactions 14: 595-604 <https://doi.org/10.1007/s11829-020-09771-8>.

Piñero, J.C., Shapiro-Ilan, D., Cooley, D.R., Tuttle, A.F., Eaton, Drohan, P., Leahy, K., Zhang, Hancock, T., Wallingford, A.K., and Leskey, T.C. 2020. Toward the integration of an attract-and-kill approach with entomopathogenic nematodes to control multiple life stages of plum curculio (Coleoptera: Curculionidae) Insects 11(6), 375; <https://doi.org/10.3390/insects11060375>.

Piñero, J.C., Souder, S.K., and Vargas, R.I. 2020. Synergistic and additive interactions among components of food-based baits underlie female fruit fly (Diptera: Tephritidae) attraction. Entomologia Experimentalis et Applicata 168: 339-348 <https://doi.org/10.1111/eea.12890>.

Bolton, L.G., Piñero, J.C., and Barrett, B.A. 2019. Electrophysiological and behavioral responses of *Drosophila suzukii* (Diptera: Drosophilidae) towards the leaf volatile β -cyclocitral and selected fruit-ripening volatiles. Environmental Entomology 48: 1049-1055 <https://doi.org/10.1093/ee/nvz092>.

Piñero, J.C., Barrett, B.A., Bolton, L.G., and Follett, P.A. 2019. β -cyclocitral synergizes the response of adult *Drosophila suzukii* (Diptera: Drosophilidae) to fruit juices and isoamyl acetate in a sex-dependent manner. Scientific Reports 9:10574. <https://doi.org/10.1038/s41598-019-47081-z>.

Piñero, J.C., Shapiro-Ilan, D., Cooley, D.R., Tuttle, A.F., Eaton, Drohan, P., Leahy, K., Zhang, Hancock, T., Wallingford, A.K., and Leskey, T.C. 2020. Controlling plum curculio adults and larvae using odor-baited trap

trees and entomopathogenic nematodes: Results from a six-year study. Fruit Notes 85(3): 1-5.

Piñero, J.C., Cooley, D.R., Clements, J., Schloemann, S., and Garofalo, E. 2020. Massachusetts Fruit IPM Report for 2019. Fruit Notes 85: 1-8.

Piñero, J.C., Cooley, D.R., Greene, D., Clements, J., Garofalo, E., Schloemann, S., Leahy, K., and Simisky, T. 2020. 28th Annual March Message to Massachusetts Tree Fruit Growers. University of Massachusetts Extension. Available at: https://ag.umass.edu/sites/ag.umass.edu/files/pdf-doc-ppt/28th_annual_march_message_2020_.pdf

Piñero, J.C. 2020. Assessment of a Non-pheromonal Lure System for Attracting Adult Tortricid Moths. Fruit Notes 85: 11-14.

Research and Extension Grants

Supporting IPM on diverse Massachusetts farms through the integration of applied research and Extension. 09/01/2017 - 08/31/2021. USDA NIFA Award Number 2017-7006-27137. H. Sandler (PI) and J. Clements (Co-PI). \$1,156,009.00.

Cooley, D. R. and J. M. Clements. Using Computer Vision to Improve Data Input for Precision Thinning Models in Apples. USDA/NIFA and NSF CPS: Medium: Collaborative Research. 6/1/2020 – 5/31/2023. \$430,762. In collaboration with Carnegie Mellon University. Total Award both Institutions \$1,100,000.

Precision crop load management of apples. USDA-NIFA-SCRI SREP 2020-51181-32197. 09/01/2017 - 08/31/2021. T. Robinson (PD), L. Chang (Co-PD), Cornell University., and J. Clements (Co-PI). \$4,800,00 total award.

Piñero, J.C. Evaluation of a grower-friendly attract-and-kill strategy for apple maggot control in New England apple orchards. New England Tree Fruit Growers Research Committee 6/1/20 – 11/30/20) \$2,600.

Multi-cultivar grafting: a new low-cost, grower-friendly Integrated Pest Management approach for key apple pests in Massachusetts. Massachusetts Society for Promoting Agriculture \$21,964 (2020-2021).



SUMMIT TREE SALES, INC.
REPRESENTING THE COUNTRY'S FINEST NURSERIES

Your One-Stop Source **For Commercial Fruit Trees**

- *New varieties*
- *Reliable rootstocks*
- *Top quality nursery trees*
- *Decades of industry knowledge*



Matthew Schuld



Jan Melvin



Dawn Melvin

**Your time is valuable,
make Summit your first call!**

800.424.2765

www.summittreesales.com •  Like us on Facebook

Early-season Performance of Diluted Concord Grape Juice and Commercial Lures at Attracting Spotted Wing Drosophila and Effects of Juice Fermentation on Trap Captures

Jaime C. Piñero and Heriberto Godoy-Hernandez

Stockbridge School of Agriculture, University of Massachusetts

Commercial synthetic food-based lures are available to monitor spotted wing drosophila (SWD). While effective, most commercial lures are based on fermentation materials and consequently they attract a comparatively high number of non-target insects. Captures of unwanted insects hinders trap performance and increases sorting time. For the past two years we have been evaluating the attractiveness of diluted Concord grape juice, a low-cost and readily available material, to male and female SWD. When diluted at a ratio of 1:3 (= 1 part of grape juice and three parts of water), diluted grape juice showed to be three times more attractive to males and females than one commercial lure under field conditions (see Piñero and Foley, 2018). In addition, grape juice diluted at the 1:3 ratio attracted significantly fewer (about three times less) non-targets than one commercial lure, highlighting a potential greater selectivity of diluted Concord grape juice.

Here, we assessed the efficacy of traps baited with diluted Concord grape juice at monitoring SWD early in the season when compared to four commercial lures. A secondary objective was to elucidate the effects of grape juice fermentation, as influenced by aging, on SWD captures in traps. For all experiments, we assessed the selectivity of the materials evaluated by relating captures of SWD to non-target insects.

Materials & Methods

Experiment 1: Early-season SWD monitoring

using grape juice and commercial lures. Here, we quantified the ability of 1-quart traps baited with diluted grape juice at the 1:3 dilution ratio (thereafter referred to as ‘DGJ’) to detect the first adult SWD of the growing season, compared with 1-quart traps baited with commercial lures. Five olfactory treatments were evaluated (1) DGJ, (2) Scentry® lure, (3) AlphaScents® lure, (4) Trécé broad spectrum PEEL-PAK multi-component lure, and (5) Trécé high-selectivity 3-component lure. DGJ-baited traps received 6 oz. of this material whereas traps baited with commercial lures received 6 oz. of unscented soapy water as a drowning solution. On May 4, five traps (one trap of each treatment) were deployed at each of four locations: Deerfield (on cherry trees), Belchertown (cherry), Whately (raspberry), and Amherst (elderberry). Traps were deployed on May 4 and they were removed on July 8, 2020. Once a week, all traps were serviced and all captured insects were brought back to the laboratory, for identification. At each trap inspection session, DGJ was replaced. The commercial lures remained in place until traps were removed.

Experiment 2. Effects of DGJ fermentation. This field test was conducted during June and July, 2019, in a cherry block at the UMass Cold Spring Orchard. The objective of this evaluation was to quantify the response of male and female SWD to volatiles emitted by DGJ (1:3 ratio) that had been aged for either, 7 or 14 days, compared to fresh DGJ. The four treatments evaluated were (1) fresh DGJ, (2) 7-day old DGJ, (3) 14-days old

DGJ, and (4) water as control. DGJ was prepared in the laboratory using 1-gallon containers with a gauze secured with rubber bands to allow for air exchange. GJD was prepared three times a week, and aged inside a chamber at 77°F and 65-75% relative humidity. The baits were prepared following a schedule that allowed for simultaneous evaluations of aged materials. Each treatment was replicated 5 times (= 5 cherry trees). Twice a week, traps were serviced, and insects were collected and transported to the lab, for identification. At each inspection session, clean traps were re-baited with newly-prepared materials.

Results

Experiment 1: Early-season comparison of diluted grape juice and commercial lures. As shown in Table 1, traps baited with diluted grape juice (DGJ) and with Trécé lures captured the first adult SWD of the season. This occurred on May 25, 2020. On June 1, DGJ was the only treatment that attracted SWD to traps. On June 10, all treatments except for AlphaScents attracted SWD. Results from the June 17 trapping date showed that the Scentry lure attracted the highest number of SWD. The Trécé broad spectrum lure ranked second on this date. DGJ attracted the highest number of SWD on June 30 and on July 8 (Table 1).

The selectivity of each treatment is presented in Table 1 (see section ‘Ratio (other drosophilids to SWD)’). DGJ was the most selective treatment (lower values

Table 1. Early-season captures (expressed as average number per trap) of spotted wing drosophila (SWD) and non-target insects (other insects belonging to the SWD family) in traps baited with diluted grape juice (DGJ) and four commercial lures at four Massachusetts locations. Boxes highlighted in orange indicate the date of first SWD captures in 2020.

Treatment	May 11	May 18	May 25	June 1	June 10	June 17	June 22	June 30	July 8
SWD									
DGJ	0	0	0.7	0.3	0.8	4.8	1.3	8.3	16.8
AlphaScents	0	0	0	0	0	4.3	2.8	2.3	2.8
Scentry	0	0	0	0	2.0	11.3	3	3.3	1.5
Trécé broad spectrum	0	0	0.7	0	0.5	8.3	1.3	3.0	1.5
Trécé selective	0	0	1.0	0	0.3	2.0	0.5	2.0	0.3
Other drosophilids									
DGJ	0	0.7	16.7	11	3.3	6.8	1.8	14.3	32.3
AlphaScents	14.3	39.7	4	6.5	18.8	58.3	23.3	8.0	13.0
Scentry	2	27	66	11.8	5	35.5	64.5	6.0	12.0
Trécé broad spectrum	55.7	88.3	76	14	54	106.8	10.5	10.3	22.0
Trécé selective	3	9.7	30.3	4	5.3	9.0	1.7	2.0	3.8
Ratio* (other drosophilids to SWD)									
DGJ	---	---	23.9	36.7	4.1	1.4	1.4	1.7	1.9
AlphaScents	---	---	---	---	---	13.6	8.3	3.5	4.6
Scentry	---	---	---	---	2.5	3.1	21.5	1.8	8.0
Trécé broad spectrum	---	---	108.6	---	108.0	12.9	8.1	3.4	14.7
Trécé selective	---	---	30.3	---	17.7	4.5	3.4	1.0	12.7

*=Numbers represent ratio of non-targets captured for each SWD captured; hence, a value of 1 would indicate one SWD captured for each non-target insect (= high selectivity). Conversely, high values indicate less selectivity for SWD. For each week, the most selective treatment (= the one with the lowest ratio) is highlighted in green.




Figure 1. Early-season (11 May to 8 July, 2020) captures of male and female spotted wing drosophila (SWD) in traps baited with diluted grape juice (DGJ) and four commercial lures, in four Massachusetts locations. Bars superscribed by the same letter are not significantly different at odds of 19:1.

represent greater number of SWD captured relative to non-targets) in 5 out of the 6 dates for which such ratios were calculated. On June 1, the only treatment that attracted SWD was DGJ, therefore no comparisons of ratios could be made. The Scentry lure had the best selectivity on the Jun 10 trapping date.

Figure 1 shows that, across the entire trapping period (May 11 to July 8) DGJ was a very attractive material to adult SWD. More specifically, DGJ captured 5.5, 2.2, 1.6, and 2.7 times more SWD than did Trécé -selective, Trécé -broad spectrum, Scentry, and AlphaScents lures, respectively (Figure 1). It is important to note that DGJ was replaced every week whereas the commercial lures were not replaced throughout the study. The expected lure longevity is 4-6 weeks for the two Trécé lures and for the Scentry lure, and 8 weeks for the AlphaScents lure.

Experiment 2: Effect of grape juice fermentation on SWD attraction. Aging DGJ for 1 or 2 weeks did no influence captures of male SWD in traps. In contrast, twice as many females were captured when DGJ was fermented for 1 or 2 weeks compared to fresh DGJ (Figure 2A). No statistical difference in captures was noted between the two levels of fermentation. Captures of non-target insects followed the same pattern




Figure 2. Captures (average number per trap) of (A) adult SWD (males and females combined) and (B) non-target insects (other insects belonging to the SWD family), according to treatment. This study was conducted at the UMass Cold Spring Orchard in 2019. For each SWD gender (top chart) and for non-targets (bottom chart), bars superscribed by the same letter are not significantly different at odds of 19:1.

described for female SWD. For instance, the number of other drosophilids was 70% and 74% greater when DGJ was fermented for 1 or 2 weeks, respectively, when compared to fresh DGJ (Figure 2B).

Conclusions

The results from the first study confirmed that DGJ is attractive to male and female SWD, it has high selectivity, and therefore this low-cost material can be used for monitoring SWD populations. Our fermentation study revealed that female captures can be increased if traps are left for up to two weeks in the field (we did not evaluate longer intervals). However, the fermentation process will also attract more non-target insects. Further research will be conducted to determine ways in which fermentation can be manipulated so as to attract fewer non-target insects without affecting SWD captures.

Acknowledgments

We thank Tim Nourse, and Tom and Ben Clark for allowing us to work on their orchards. We also thank Emily Begonis for assistance. Funding for this research was provided by the Massachusetts Department of Agricultural Resources Specialty Crops Block Grant with funds provided by the United States Department of Agriculture (USDA), Agricultural Marketing Service through agreement number AM180100XXXXG025. The views or findings presented here are the Grantee's and do not necessarily represent those of the State or the Missouri Department of Agriculture.

References

Piñero, J.C. Foley, N. 2018. Evaluation of diluted grape juice as an inexpensive attractant for the invasive fruit pest spotted wing *Drosophila*. *Fruit Notes* 83(3) 1-7.

LEADER OF THE PAK

REARS

THE MOST POPULAR 3 PT. HITCH WE SELL!

REARS PAK-TANK SPRAYERS

- Stainless steel tanks, mechanical agitation
- PTO shaft driven diaphragm pump
- Variety of booms, guns and hose available
- Excellent for small fruit, Christmas trees and general row crop spraying

**25 TO 200 GALLON SIZE • 3 PT. HITCH OR FRONT MOUNT
STANDARD OR HEAVY DUTY FRAMES**



100 gallon Pak-Tank



Call for a catalog or just stop by

You will always be able to speak to a knowledgeable, friendly person who can help.

www.oescoinc.com

800-634-5557

Mon. - Fri. 7 a.m. - 5 p.m. & Sat. 7 a.m. - noon

8 Ashfield Rd./Rt. 116, P.O. Box 540, Conway, MA 01341

EVERY FIELD

EVERY ACRE.

EVERY ROW.



PUSH THE LIMITS

***Servicing the New Jersey Horticulture Industry with
expertise in fertility and micronutrient programs and crop
protection recommendations***

Bloomsbury, NJ
908-479-4500



GROWMARK FS





THE LEADING SMALL FRUIT NURSERY TODAY!

NOURSE

The Best Berry Plants Since 1932

- *Excellent Customer Service*
- *Wide Variety Selection*
- *Technical Support*
- *Complete Lab Facility
for Tissue Culture
& Virus Indexing*



Strawberries Currants
Brambles Gooseberries
Asparagus Elderberries
Blueberries Rhubarb

NOURSE FARMS, INC
41 RIVER ROAD
SOUTH DEERFIELD MA 01373
413.665.2658
INFO@NOURSEFARMS.COM NOURSEFARMS.COM

Eco-Friendly Insect, Disease, Bird Control

University/USDA tested

Stink Bug Traps

Brown Marmorated and Native Bugs

Insect Traps and Lures

*Plum Curculio Trap Tree **Control**,
Codling & Oriental Moth, Cranberry
Pests, **Black Stem Borer**, Others*

Honey Bee Lure

Attract Bees - Increase Pollination

Predalure attracts beneficials

Oriental Beetle MD

*Mating Disruption
Fruit Crops & Ornamentals*

Prestop

*New Biofungicide Impressive
Activity. Foliar/Root Diseases*

Avex

*Bird Control. Apply by ground or
air. Cherries, Blueberries, Sweet
Corn, other crops*



*Committed to the Environment and Green Technology
Since 1990*

P. 303-469-9221
agbio@agbio-inc.com
www.AgBio-Inc.com



Thwarting the Low-spray Plan: Marssonina Throws a Curve Ball

Elizabeth Garofalo and Daniel Cooley
University of Massachusetts

Marssonina coronaria, the fungal organism known to cause Marssonina leaf blotch (MLB) on apple, is an emerging disease in the Northeast. The pathogen was first documented in the United States in 1903 (Davis 1903). At the time, it was listed under a different name: *Ascochyta coronaria*, and has since undergone several changes in its taxonomic name before arriving at today's *M. coronaria*. Until recently, the disease has been considered minor in the US, compared to other diseases caused by fungi, notably apple scab, powdery mildew, rusts and fruit rots.

Over the last 20 years, *M. coronaria* has become a commercially significant pathogen in organic apple growing regions in Europe. It caused significant damage on 'sustainably grown' apples in northern Italy in 2002 and 2003 (Tamiatti and Matta 2003). It now causes losses in southern Germany, southern Austria and Switzerland as well. (Persen et al. 2012; Wöhner and Emeriewen 2019). In China, India and other Asian countries, MLB has caused huge losses over the past 30 years, and is considered one of the most important diseases in apple production over that region (Verma and Sharma 2004; Lee et al. 2011; Dang et al. 2017).

M. coronaria has been observed in commercial orchards in the north-eastern United States in recent years, and is becoming especially problematic in organic and low-spray orchards. It is also being seen in conventional orchards. Symptoms first begin as discreet dark brownish-purple spots. These spots develop acervuli, a fungal fruiting body that produces spores (Figure 1). These spots expand into yellowed chlorotic regions which then coalesce into a mottled yellow "blotch" of the leaf. If infections are

severe enough, trees defoliate prematurely. Over a few years, this can lead to a decline in tree health, and cause a reduction in fruit production and quality. *M. coronaria* can also cause spots on fruit, decreasing value or even making them largely unmarketable. Fruit symptoms are less common than foliar symptoms.

Currently, there are no fungicides labeled in the United States for use in managing *M. coronaria*. Other regions that have been struggling with MLB in commercial orchards report that there are materials that are effective in preventing infection. For example, in India broad-spectrum protectant fungicides including mancozeb, ziram, dodine, chlorothalonil and thiophanate-methyl controlled MLB, but sterol inhibitors (DMIs) were not effective (Verma and Sharma 2003). Dang




Figure 1. Leaf infected with Marssonina leaf blotch. Red arrows point to magnified acervuli, which are still difficult to see but are a good field diagnostic characteristic.

et. al. (2017) in China tested the strobilurin kresoxim-methyl (Sovran), DMI fungicides (tebuconazole = Tebuzole, myclobutanil = Rally, difenoconazole = one a.i. in Inspire Super, and two others not sold in the US), and Bordeaux mix. Sovran was effective, but performance of the DMIs and Bordeaux were erratic, with tank-mixes of different DMIs or with Bordeaux more effective. This work also recommended an early summer application of mancozeb in addition to DMI fungicides is effective in preventing disease in the field.

A trial looking at fungicide performance against apple fruit rots in Pennsylvania also proved useful in evaluating fungicide efficacy against MLB, and showed that several fungicides were very effective against the

personal communication).

Selecting disease resistant apple varieties might mitigate the impact of *M. coronaria*. Little is known about the relative susceptibility of apple cultivars to MLB, including the susceptibility of cultivars resistant to scab, and in some cases, other diseases. Commercially available cultivars selected for disease resistance were not screened for tolerance to MLB, as they were bred well before *M. coronaria* was observed on commercial apple trees in the U.S. Additionally, recent breeding has focused less on disease resistance and more on consumer sensory appeal (Peace et. al. 2019). In order to manage MLB, it is important to determine which cultivars currently on the market, or in the ground, are resistant to *M. coronaria*. In this article, two scab-resistant apple cultivars, GoldRush and Enterprise, grown under a low-spray, organic program are assessed for susceptibility to MLB during the 2020 season.

GoldRush is a commercially available scab resistant apple variety bred by the Purdue-Rutgers-Illinois (PRI) apple disease resistant breeding program (Janick 2006). In addition to field immunity to apple scab, GoldRush exhibits high resistance to powdery mildew and is moderately resistant to fireblight. It also stores extremely well, (Crosby et. al. 1994a). Goldrush develops optimum flavor/sugars after 4-6 weeks in cold storage. It maintains its flavor and texture for 10-11 months. Quality



Figure 2. Enterprise (foreground) exhibits resistance to Marssonina compared to GoldRush (background).

disease if applied from pink through the summer (K. Peter). Captan, mancozeb, Luna Sensation, Luna Tranquility and Merivon all provided excellent control. Topsin, Flint Extra, sulfur and the SDHI fungicides (Aprovia, Fontelis and others) gave good to excellent control. Again, these materials are not labeled specifically for use against MLB in the US. However, they may be used to manage other summer diseases such as sooty blotch/flyspeck, bitter rot and black rot. Mancozeb applications made early in the apple growing season targeting apple scab and rusts can also manage early season *M. coronaria* infections in conventional orchards. But mancozeb has a 77-day preharvest interval, so other fungicides or methods need to be used during the summer against MLB. For organic growers, the situation is difficult, as organically approved fungicides, except for sulfur, are not very effective against MLB (K. Peter,

can be enhanced for the longer storage times if treated with Smartfresh (personal communication, Win Cowgill, Professor Emeritus, Rutgers Univ.). Goldrush makes excellent hard cider as tested by Duane Green in his UMass hard cider apple trial. Is is utilized commercially for hard cider production both in NJ and PA cideries. It is also planted extensively in NJ, PA, and OH, as it has developed a retail sales and PYO following due to its high desert quality. Goldrush ripens 3-3.5 weeks after Delicious and after Fuji but before Pink Lady. Because of its long maturity development and late harvest, there was concern by northern growers that it would not ripen in some years. Dr. Ian Merwin found that several nights of below 32F will trigger ripening, and it could be harested in good condition. Because of this, Goldrush can usually be harvested in Geneva, NY before it is ready for harvest in Norhtern

NJ. (personal communication, Ian Merwin, Professor Emeritus, Cornell Univ.). Generally, GoldRush is not prone to premature drop.


Enterprise is another commercially-available apple variety developed in the PRI program. This late maturing, approximately three weeks after 'Delicious', apple. Enterprise was extensively tested in greenhouse conditions and proved resistant to apple scab. It is also highly resistant to cedar apple rust and fireblight and moderately resistant to powdery mildew. This apple is reported to keep well, maintaining "excellent" quality up to six months in storage. Enterprise has a severe calcium disorder "Mystery Spot" that limits its commercial acceptance. (Tietjen et.al 1994, Tietjen et. al.1995). While Enterprise has excellent flavor and large size, it also has tough skin. These two issues limit it from being adopted as a retail apple cultivar (Cowgill et. al. 1994). Note that Enterprise is one of the parents of Washington States new club variety Cosmic Crisp tested as 'WA 38', the other parent being Honeycrisp. Of interest is that after 2 years of commercial production Cosmic Crisp is having a calcium disorder problem they coined 'green spot' (Courtney, 2018).

The original Enterprise seedling was planted out in 1982 at the Purdue University Horticultural Farm. The first GoldRush seedling was planted in 1973(Crosby et. al. 1994b). Both cultivars attribute their apple scab resistance to the V_f gene derived from *Malus floribunda* parentage.

Differences in the parentage of Enterprise and Goldrush suggest there may be differences in MLB susceptibility. Two varieties common in the parentage of these varieties have shown susceptibility to MLB, Golden Delicious and Rome Beauty (Li et al. 2012; K. Peter, personal communication, October 9, 2020). Golden Delicious is a direct parent of GoldRush. While Enterprise also has both Golden Delicious and Rome Beauty in its background, it is a direct result of a "sibling cross", where two plants

grown from seed of the same fruit are bred together to create the new cultivar, in this case, Enterprise. Enterprise is three generations removed from Golden Delicious and four from Rome beauty. GoldRush has no generational separation from Golden Delicious though it is five generations removed from Rome Beauty, but with fewer intervening crosses than Enterprise. These sibling and other intervening crosses in the parentage of Enterprise as compared to GoldRush make it less directly descended from cultivars known to be susceptible to MLB, suggesting it may be less susceptible to MLB. It appears as though the gene deck is stacked against GoldRush when it comes to Marssonina.

In the comparison of MLB susceptibility of GoldRush and Enterprise, a rating scale of 0 to 9 was used to quantify disease severity, with 0 having "no visible lesions", and 9 showing "tree almost completely bare, only few leaves left on top or side branches" (Bohr et. al. 2018). Four trees were selected at random from each of six rows from each cultivar providing 24 trees as-



sessed per cultivar and a total of 48 trees assessed. All trees assessed had symptoms of MLB. The severity of MLB on GoldRush was significantly higher than that of Enterprise (Figure 1). MLB can also cause premature fruit drop in GoldRush, which appears to be more susceptible than Enterprise (Figure 2).

MLB apparently is becoming established in our region, and when traditional fungicide applications are stretched, or in the case of some low-spray or organic orchards, eliminated altogether, significant damage can occur. As MLB becomes more prevalent it is increasingly important to understand which varieties are more or less susceptible. Some evaluations of commercial cultivars in China (Li et al. 2012) show there are significant differences. Wöhner and Emeriewen (2019) identified differences in cultivar susceptibility drawing on several sources in their review of MLB. However, they rate Honeycrisp as resistant, while experience in the Northeast indicates it is quite susceptible. More research on resistance is needed. Unfortunately, it appears that available disease resistant varieties may not fare any better than many commercial varieties against MLB. Identifying which varieties are tolerant and which susceptible to the disease is a critical first step in managing this emerging disease.

Literature Cited

- Bohr, A., Buchleither, S., Hechinger, M. and Mayr, U. et al. (2018). Symptom occurrence and disease management of Marssonina blotch. 18th International Conference on Organic Fruit-Growing: Proceedings of the Conference, 19-21 February 2018, Hohenheim, Germany.
- Courtney, Ross. 2018. Cosmic conundrum: What's causing these spots? GoodFruit Grower. <https://www.goodfruit.com/cosmic-conundrum-whats-causing-these-spots/>
- Cowgill, W.P., Jr., M.H. Maletta, W.H. Tietjen, J. Compton, D.F. Polk, J.C. Goffreda. 1994. "Preliminary Performance of Six Scab Resistant Apple Cultivars in Northwest New Jersey." Proceedings 70th Cumberland-Shenandoah Fruit Workers Conference, Nov. 17-18, 1994, Horticulture.
- Crosby, J. A. Janick, J. Pecknold, P. Joffreda and J. Korban, S. (1994a). 'GoldRush' Apple. HORTSCIENCE 29(7):827-828.
- Crosby, J. A. Janick, J. Pecknold, P. Joffreda and J. Korban, S. (1994b). 'Enterprise' Apple. HORTSCIENCE 29(7):825-826.
- Dang, J. L., Gleason, M. L., Niu, C. K., Liu, X., Guo, Y. Z., Zhang, R., et al. 2016. Effects of Fungicides and Spray Application Interval on Controlling Marssonina Blotch of Apple in the Loess Plateau Region of China. Plant Dis. 101:568-575.
- Davis, J. J. (1903). Third supplementary list of parasitic fungi of Wisconsin. Transaction of the Wisconsin Academy of Science, Art and Letters, 14(1), 83-106.
- Janick, J., 2006. The PRI apple breeding program. HortScience, 41(1), pp.8-10.
- Lee, D.H., Back, C.G., Win, N.K.K. & Choi, K.H. 2011 Biological characterization of Marssonina coronaria associated with apple blotch disease Mycobiology 39 200 205.
- Li, Y., Hirst, P.M., Wan, Y., Liu, Y., Zhou, Q., Gao, H., Guo, Y., Zhao, Z., Wang, L. and Han, M., 2012. Resistance to Marssonina coronaria and Alternaria alternata apple pathotype in the major apple cultivars and rootstocks used in China. HortScience, 47(9), pp.1241-1244.
- Peace, C. P. Bianco, L. Troggio, M. Van de Weg, E. Howard, N.P. Cornille, A. Durel, C.E. Myles, S. Migicovsky, Z. Schaffer, R.J. Costes, E. Fazio, G. Yamane, H. Van Nocker, S. Gottschalk, C. Costa, F. Chagné, D. Zhang, X. Patocchi, A. Gardiner, S. E. Hardner, C. Kumar, S. Laurens, F. Bucher, E. Main, D. Jung, S. and Vanderzande, S. (2019). Apple Whole Genome Sequences: Recent Advances and New Prospects. Horticultural Research. 6:59
- Persen, U., Steffek, R., Freiding, C., & Bedlan, G. (2012). Erstnachweis von *Diplocarpon mali* an *Malus domestica* in Österreich. Journal für Kulturpflanzen, 64(5), 168-170.
- Tamietti, G. and Matta, A., 2003. First report of leaf blotch caused by Marssonina coronaria on apple in Italy. Plant Disease, 87(8), pp.1005-1005.
- Tietjen, W. H., W.P. Cowgill, Jr., K.S. Petersen, D.F. Polk, G. Slifer. 1995. "The Effect of Calcium Sprays on Incidence of 'Mystery Spot' on Enterprise Apple." Proceedings 71st Cumberland-Shenandoah Fruit Workers Conference.
- Tietjen, W.H., D. F. Polk, W.P. Cowgill, Jr., K.S. Petersen, E.F. Rizio, Jr. 1994. "Mystery Spot of 'Enterprise' Apple." Proceedings 70th Cumberland-Shenandoah Fruit Workers Conference, Nov. 17-18, 1994, Horticulture.
- Verma, K.D. and Sharma, J.N., 2004. Scab and Marssonina Blotch –Threatening Diseases of Apple and Their Management. Annual Review of Plant Pathology (Vol. 2), 2, pp.207-220.
- Wöhner, T. and Emeriewen, O.F., 2019. Apple blotch disease (Marssonina coronaria (Ellis & Davis) Davis)—review and research prospects. European journal of plant pathology, 153(3), pp.657-669.



Adams County Nursery, Inc.

Growers of Fruit Tree Nursery Stock

26 Nursery Road, Aspers, PA 17304

acn@acnursery.com • 717.677.8105 • www.acnursery.com



Northern New Jersey Report, As Presented at the NEIPM Conference on October 2020

Win Cowgill

Professor Emeritus Rutgers University, Win Enterprises International, LLC

Spring Weather


It was a tough spring for New Jersey growers. We started the season almost 3 weeks early with green-tip in Hunterdon County on March 16. While peaches were early as well, they have managed to move through bloom. Apples dragged along and were at tight cluster to early bloom on April 28 on many varieties depending on location and orchard exposure.

In North Jersey, nine freeze events occurred: 4/17, 4/18, 4/19 and 4/20, 4/22, 4/23, 5/8, 5/12, 5/13. Some apples were hurt in southern NJ. Cold events ranged from 32F down to 23F, but not every farm had all

events. Bloom and post bloom damage occurred at most orchards, some severe some minimal, some lost whole crops, apples and or peaches. Damage depended on location, cultivar, stage of bloom, etc. Three types of damage was observed: ovary, stigma (where pollen germinates) and style, anther (pollen)- & filament. Some had all three areas injured. Overall, I gauged the apple crop at 60% of a normal crop in NJ.

Apple Thinning

Fruit set and thinning were a challenge this year with these cold events. I had several growers use Pro-






Figure 2. Rimpro Fireblight (*Erwinia*) Baptistown, NJ for 2020. Fireblight infections reached critical levels on several dates according to Rimpro but none reached the infection threshold of 0.2. The Critical fireblight potential infection of May 16 did not reach the infection potential of 0.2 but on NEWA (see Figure 3) it reached 113 EIP which is the minimum for infection. Growers with open bloom should have sprayed on the May 15-17 with Strep.

malin or Perlan to help set apple fruit during several of the most severe events, with some success. We also enhanced typyness on many varieties, like Gala, with the Promalin treatment. A few growers on a few cultivars needed to bloom thin, but many held off. Where fruit set appeared strong, we thinned at petal fall and often applied a second 8-14 mm spray. Many growers with rapidly sizing fruit felt they needed to thin again at 14-18mm. The Cornell Carbohydrate Thinning Model showed a severe carbohydrate deficit during this time period. Even though we were conservative with this spray, some blocks thinning more than desired (overthinned), resulting in some reduced crop.

Diseases

Apple Scab and Fireblight were monitored with NEWA (<http://newa.cornell.edu/>) (Milford, NJ) and Rimpro (<https://www.rimpro.eu/>) at my location (Baptistown, NJ). Rimpro is a paid subscription service at your location using either your own weather station data or using virtual weather forecasting via Metro Blue. NJ maintains a set of weather stations in NJ through the Rutgers NJ weather network (<https://www.njweather.org/>), and NJEAES/Rutgers Coop. Extension

pays Cornell Subscription fee. You also can have your own weather station and pay NEWA a subscription fee directly to Cornell for connection.

Fireblight was not a severe issue in Northern NJ in 2020. Several fireblight infection events occurred during bloom. Growers that used the program of applying low rates of Apogee beginning at pink, through first cover fared best with the least amount of infection. Note that I work with growers in 5 states. One grower in Orange County, NY had a severe fireblight outbreak in a newly planted block of mixed apple varieties. The Rimpro and NEWA models showed close to 20 days of high susceptibility. Several streptomycin sprays did not control the infection, since they were not started soon enough, and not enough applications were made. It appears that fireblight may have come in on the susceptible M.9 Nic 29 rootstock from an out-of-state nursery. Tested by Cornell it was not a streptomycin-resistant strain. Late-blooming, newly-planted apple trees are always at risk for fireblight and must have the flowers removed before blooms open or be covered with streptomycin 24 hours before and after infections. A low-dose apogee program would also be helpful.

We are working on a new gibberellin formulation that will be used to treat trees in the production nursery to eliminate bloom of trees when they are planted in the orchard, this will be a big boon to apple growers. It looks very promising.


Brown Marmorated Stink Bug

We had several large populations of brown marmorated stink bugs (BMSB) in Northern NJ in late August through September that apple growers needed to treat. Rutgers has NOT been obtaining section 18's on key insecticides (Including Bifenthrin) for the past 3 years, so options were limited for control. Fortunately, thanks to Peter Jentsch/Cornell, we had the use information and the newly labeled Venerate XC biological insecticide that works on BMSB, and other insects to keep them from feeding on fruit. Several growers were able to applied Venerate XC to several large Honeycrisp blocks with good control. From Peter Jentsch -Cornell Blog: <http://blogs.cornell.edu/jentsch/2020/08/24/management-options-for-the-stink-bug-complex-on-pome-fruit-near-harvest-august-24th-2020/>.

Venerate XC is a novel approach. Keeping BMSB from feeding on the fruit is at the crux of our stink bug management program as we near harvest. Killing the insect as it moves into the orchard and onto the fruit is the traditional method of crop protection and proven to be very effective. Yet limitations, based on the material pre-harvest intervals constrain their use near harvest. There are no effective pyrethroids, neonicotinoids or pre-mixed insecticide tools that permit their use within the 7-day to harvest window with very few options available during the days prior to fruit harvest. The development of newer classes of insecticides that produce an anti-feeding response in the pest provides an additional mode of action for BMSB management. Both Corteva insecticide Closer 240 SC (classified

as a Group 4C insecticide / neonicotinoid – 7 days to harvest) and Venerate XC (microbial-based insecticide with multiple modes of action and 0 days to harvest) have anti-feeding activity against BMSB. The 0 days to harvest of Venerate provides BMSB management up to the day of harvest under high risk conditions. Peter's complete study on Venerate Xc can be found in his 2017 report, <https://cpb-us-e1.wpmucdn.com/blogs.cornell.edu/dist/f/3191/files/2013/10/2017-Final-Report.8.14.18-pnm7v2.pdf>.

The Venerate XC label requires a 4 hr. REI and has a 0 days PHI. In mixed blocks often requiring multiple picks, the use of Venerate XC has been shown to effectively reduce BMSB feeding injury to fruit 7 days prior to harvest. Although this insecticide provides no toxicity to the insect, it effectively reduced feeding over 7 days. A Technical Information Bulletin is available for the use of Venerate XC bio-insecticide brown marmorated stink bug on apple.



New Jersey News

Melick's Town Farm Adds New Cold Storage and Production Building

Win Cowgill

Professor Emeritus Rutgers University, Win Enterprises International, LLC

Melick's Town Farm has a long-planned construction project under way to add cold storage and production space to the Oldwick operation. See photos. Melick's is the largest apple grower in New Jersey with 25,000 apple trees and 5,000 peach, nectarine, plum, and pear trees. Their orchards cover 120 acres of their 650 acres in Hunterdon County, New Jersey. <https://www.melickstownfarm.com/>.

The farm is run by 10th generation farmers, siblings Peter, Rebecca, and John Melick with the continued support of their parents George and Norma.

John Melick indicated that "we needed more cold storage space for the past 25 years and this new addition will meet our



needs now and for the future." The building is designed to have cold storage for 1000 apple bins plus additional cold storage for the hard cider products. In addition, the building will be equally divided between production/grading/staging space, a new hard cider production facility, and cold storage.

Photo credits: John Melick.

William J. Roberts, World Renowned Rutgers University Agricultural Engineer Passed May 21, 2020

William J. Roberts, a lifelong resident of New Monmouth, passed away on May 21 at the age of 88.

Bill was famous for his invention of the air-inflated double-layer polyethylene greenhouse. This greenhouse technology revolutionized production and was adopted around the world. The impact of his life's work is amazing.

In 1999, there were approximately 23,125 acres of air-inflated double-layer polyethylene greenhouses in production in the USA and 1,705,125 acres worldwide. Approximately 65% of all commercial greenhouses in the United States use the air-inflated system.

He was born on his family's farm to the late Thomas S. and Helen (Conover) Roberts. At a Rutgers Cook College retire luncheon January 2020 I got to visit with my friend Bill Roberts for 15 minutes or so. One of the stories he shared was that he still lived on the family farm. He said he could look out his window at the original farmhouse and see the window to the bedroom he was born in.

Bill graduated from Leonardo High School in 1949 and earned his Bachelor and Masters of Science degrees in Agricultural Engineering at Rutgers University. Having completed ROTC, he served his country from September 1953 to July 1955 serving in the Far East Command in Japan as a First Lieutenant. He then taught at Cook College, Rutgers University, where he achieved Professor II status, served as a Department Chairman for 23 years, and worked as a Professional Engineer and extension specialist in bioresource engineering. Bill retired from Rutgers in 1999 after a distinguished 41- year career.

His work and passion has left an indelible imprint across the globe in the form of the air-inflated, double-layer polyethylene greenhouse, an innovation that

revolutionized the greenhouse industry worldwide. When asked about this project, Bill would modestly say, "everyone has a good idea once in a while."

The site of the original greenhouse at Rutgers was designated a national agricultural (ASABE) historic

landmark in 2004. As a Professor Emeritus at Rutgers, he received the Distinguished Service in Agriculture Award from the New Jersey Farm Bureau, the Agribusiness Heritage Award and the Gold Medallion Award from the New Jersey Agricultural Society. He was named a Fellow for his national professional association, the American Society of Agricultural and Biological Engineers. These are only a few of the awards for his outstanding contributions to New Jersey and global agriculture. Throughout his life, Bill was an active member of New Monmouth Baptist Church and



proud to be the great-great-grandson of its founding pastor. His devotion to his Lord and Savior was evident to all who knew him. For over 65 years, he served the church in various roles.

Along with his wife, Dottie, he was a youth leader for twenty years and led a Wednesday night Bible study for over forty years until his death. Over the years, they also led mission trips to Appalachia and Ecuador and were active members of the church choir. He taught a Sunday School class beginning while in Japan and continuing on until the present as well. He was a deacon for over fifty years, exemplifying servant leadership and mentoring younger church leaders who now serve throughout the world. He will also be remembered specifically for his hugs, jokes, and stories. Bill also served as the faculty advisor for Campus Crusade at Rutgers and on multiple mission boards, including Farms International and Chapin Living Waters. His love, faithfulness, and hospitality impacted countless

lives as he strove to follow Christ. The resurrection of Christ was the foundation of all Bill did and the source of hope for his future.

Bill is predeceased by his brother and sister-in-law, John and Elizabeth Roberts. Surviving are his beloved wife of 66 years, Dorothy (Stevens) Roberts, his loving daughter and son-in-law, Deborah Roberts and James Marcantonio, his cherished granddaughter, Jillian Roberts Marcantonio, and his dear brother and sister-in-law, Thomas and Patricia Roberts. Also surviving are his nieces and nephews, Douglas Roberts and his wife Ann,

Betty Cardinale and her husband Carl, Jean Herholdt and her husband Dan, and John Roberts and his wife Lori, as well as his cousin, Gertrude Lindemann.

A private service for the family has been held at Fair View Cemetery, where Bill served as a trustee for many years. A public celebration of life will follow at a later date. Memorial donations may be sent in Bill's memory to New Monmouth Baptist Church, 4 Cherry Tree Farm Road, Middletown, NJ 07748. Checks made out to New Monmouth Baptist Church, memo line: Roberts Memorial Fund.

John “Jack” J. Kupcho Long-term Chair of the Rutgers Agricultural Agents Department Passed on May 11, 2020

John J. (Jack) Kupcho, 83, died on May 11, 2020 at the Christian Healthcare Center in Wyckoff, NJ. Private Services were held.

As Chairman of the Department of Agricultural and Resource Management Agents at Rutgers' Cook College, he was responsible for providing program leadership and staff development for 57 agents and program associates in 20 New Jersey counties.

Jack began his career as the Essex County Agricultural Agent at Rutgers Cooperative Extension in 1963, a time of great change and development in the North Jersey agriculture industry. Jack assisted commercial vegetable growers in Bergen, Essex, Morris and Passaic counties in changes to their production methods to maximize resources. Jack developed the Master Gardener Program for Northeastern New Jersey. He organized leaf composting and ground water education programs for community leaders.


Jack was active in many agricultural organizations –the Essex County Board of Agriculture, National As-

sociation of County Agricultural Agents, Agricultural Agents Association of New Jersey, Northeast Agricultural and Natural Resource program Leaders Committee, Northeast Community Resource Development Program Leaders Committee, and Northeast Center for Rural Development.

For his 51 years of service to teaching, advising and improving New Jersey agriculture, Jack received the 2015 New Jersey State Board of Agriculture's Distinguished Service to Agriculture Award.

Active locally, Jack was involved with the West Caldwell Environmental Commission, where last August he was honored by having the West Caldwell Community Gardens named after him.

Jack is survived by his wife, Helen Januszewski Sabo Kupcho; his daughters Terri Schwenker (Donald), Mary Ellen Will (Brad), Kathy Lynn Canale (Gary), Colleen Tocci (Craig) and grandchildren Madison, Lauren, Claudia, Rachel, Megan, Julia, Abby, Caitlin, Kylie, and Jack.



Fresher, Sweeter, Juicier

Peaches From Circle M Farms



*Exceptional quality and flavor at attractive prices
Brought to you from the 4th generation family
farm of Santo John Maccherone*

- PLU Stickers Available Upon Request
- Low in Calories
- Nutritious and Delicious
- Exceptional Flavor



SANTO SAYS...

"Peaches from Circle M Farms just plain taste better. Their size and color are what your customers want, and their flavor is the icing on the cake.

That's because our orchards are tended with care and attention, and their fruit is left on the tree to ripen naturally until they reach their optimum sweetness. Then they are picked by hand, taste-tested by me to ensure quality, and packed gently into

25 lb. volume boxes for shipping. Single layer cartons are available upon request.

Circle M's 30 varieties of peaches and nectarines are known for their sweetness and their luscious color. They look as good as they taste.

Discriminating customers appreciate our white flesh peaches because of their lower acid levels and supremely sweet taste.

Add it all up: Peaches from Circle M Farms are fresher and sweeter, picked by hand when their time has come, and packed with care for a longer shelf life.

Yellow, white and donut peaches and yellow and white nectarines are available early July to late September."



**Call us at 856 856 207-4542 or
609 381-6465. Fax your order to
856 878-0404**



Available in 16 oz. and 64 oz. sizes

**Made from fresh New Jersey Peaches
"Peach Cider Drink, Peach Salsa, Peach
Preserves"**

**Santo John Maccherone
circlemfarmsllc@gmail.com**

**Circle M Farms
88 Sandy Ridge Road
Salem, N.J. 08079**



Horticultural News

New Jersey State Horticultural Society
176 Airport Road
Hackettstown, NJ 07840