Monitoring Egg Parasitoids of the Brown Marmorated Stink Bug in Massachusetts

Mateo Rull-Garza, Zoe Robinson, Jaime C. Piñero Stockbridge School of Agriculture, University of Massachusetts Amherst

The growth and spread of invasive pests like the brown marmorated stink bug (BMSB) is not usually encumbered by natural enemies (e.g., predators, parasites), which are not well-suited to interact with the invasive species (Venette & Hutchison, 2021). Instead, the responsibility to control the pest falls on growers, who often rely on chemical pesticides. However, the use of broad-spectrum insecticides can kill and deter beneficial insects that naturally control BMSB population levels (Leskey et al., 2012).

In a variety of cropping systems throughout the US, there are native species of parasitoids that attack stink bug eggs (Tillman, 2010). Over the decades, several uninvited biocontrol candidates have popped up on new continents. One example is the exotic samurai wasp (SW), a parasitoid that, though smaller than a tenth of an inch, is capable of killing up to 80% of BMSB populations in Northern China (Ogburn et al., 2021). This tiny wasp achieves such powerful control by laying its eggs inside BMSB eggs, effectively killing the developing stink bug nymphs as their offspring develop. The SW was first detected in Maryland in 2014 and, as of 2022, has been found in at least 13 states, including our neighboring New York State (Ogburn et al., 2021). We do not yet know if the SW is present in Massachusetts. If present, the SW could become a *biological control* agent critical to the suppression and control of BMSB.

Here, we report the results of field surveys aimed at assessing the presence of the SW in Massachusetts and identifying the complex of BMSB egg parasitoids that already exist in the state.

Materials and Methods

Sentinel Egg Preparation. We purchased BMSB eggs from the Phillip Alampi Beneficial Insect Laboratory (New Jersey Department of Agriculture, NJ). Upon arrival, egg shipments were frozen in -80 degrees Celsius (Tillman et al., 2020) for either 48 hours (May through mid-August) or 120 hours (three shipments in late August and September). After freezing, egg masses were individually attached to a piece of white paper using a small amount of superglue and left to dry for at least 30 minutes before deployment. In the field, prepared sentinel egg masses were stapled on the

underside of leaves from trees in the orchard perimeter (Ogburn et al., 2021). Specifically, eggs were deployed on/ near plant species known to be attractive to either the BMSB or wasp parasitoids.



Figure 1. Deployment of brown marmorated stink bug masses in the field.

Study Sites. For sentinel egg deployment, we selected ten fruit tree orchards located in the outskirts of either rural or suburban centers across the state of Massachusetts based on **BMSB** habitat suitability data (Figure 1). The ten or-



chards were visited at least once per month except for May, when only four of the orchards were visited.

Incubation and assessment of parasitism and predation. After 72 hours in the field, the sentinel egg masses were retrieved and counted, noting chewing predation damage (Tillman et al., 2020). Predated eggs were discarded, and the remaining were placed inside petri dishes unique for each orchard. The tops of the petri dishes were covered in four layers of insect netting (1×0.5 mm) before being incubated. Eggs were left inside the incubator until parasitoids emerged and died, at which point the petri dishes were removed, and the parasitoid wasps were transferred into 70% ethanol and refrigerated for preservation. After five weeks and no new parasitoid emergence, the remaining eggs were dissected to recover underdeveloped parasitoids that were not able to emerge. we doubled the number of eggs deployed to 2,556 and recorded 28 eggs (1.1%) with chewing damages, while no parasitism was recorded. July was the first and only month when we recovered parasitized eggs (84), constituting a 2.5% parasitism rate out of the 3,405 eggs deployed in that same month. The adult parasitoids that actually emerged were *Trissolcus euchisti* (n= 53), *Telenomus podisi* (n= 10), and *Ooencyrtus* sp. (n= 7), as well as two underdeveloped parasitoids from the subfamily *Telenominae* (Figure 2).

We also recorded the highest levels of predation in July, when 328 eggs (9.6%) showed signs of chewing. Out of 1,888 eggs recovered in August, 71 of these (3.8%) showed signs of predation. No parasitoid wasps emerged in August. Only two orchards were visited in September and 738 eggs were deployed. Out of these, 16 (2.2%) showed signs of chewing and we recorded no parasitoid emergence (Table 1).

Results

All results are summarized in Table 1. In May, we recovered 1,224 eggs from 4 orchards, out of which 6 (0.5%) had signs of chewing predation, and none hatched wasp parasitoids. In June, **Table 1.** Summary of predation and parasitism on Brown Marmorated Stink Bug sentineleggs. Recovered egg masses were assessed to identify chewing predation damage.Parasitism was noted as parasitoids that either, emerged or were dissected out from theeggs if underdeveloped. Highest rates of predation and parasitism were observed in July.

Month	Eggs recovered	Eggs chewed	% Eggs chewed	Eggs parasitized
May	1224	6	0.5	0
June	2556	28	1.1	0
July	3405	328	9.6	84 = (2.5%)
Aug	1888	71	3.8	0
Sep	738	16	2.2	0

Conclusion

While we did not find evidence of SW presence in the Massachusetts orchards that were surveyed, across five out of the nine participating orchards we observed 2.5% parasitism of sentinel eggs and recovered 72 parasitoid wasp specimens. While parasitism rates were not substantial, few orchards in MA currently report BMSB infestation problems, which could indicate low BMSB population levels. With few BMSB eggs around, native wasps may preferentially parasitize well-established host populations. According to our data, BMSB egg predators currently seem to exert more powerful control than parasitoid wasps, as we report chewing predation rates as high as 9.6% mid-summer. Our surveillance efforts will continue in 2023.

Acknowledgements

We appreciate the support of Dr. Elijah Talamas who helped us identify

the wasp parasitoids. Thank you to Heriberto Godoy for help in the field and to all of the MA growers who agreed to participate in our study. Funding for this research was largely provided by The Massachusetts Department of Agricultural Resources (MDAR) using Specialty Crops Block Grant funds provided by the United States Department of Agriculture, Agricultural Marketing Service. The views or findings presented here are the Grantee's and do not necessarily represent those of MDAR.

Literature Cited

Leskey, T.C., Short, B.D., Butler, B.R. & Wright, S.E. (2012) Impact of the Invasive Brown Marmorated Stink Bug, Halyomorpha halys (Stål), in Mid-Atlantic Tree Fruit Orchards in the United States: Case Studies of Commercial Management. Psyche: A Journal of Entomology, https://doi.org/10.1155/2012/535062

Ogburn, E.C., Heintz-Botz, A.S., Talamas, E.J. & Walgenbach, J.F. (2021) Biological control of Halymorpha halys (Stål) (Hemiptera: Pentatomidae) in apple





orchards versus corn fields and their adjacent woody habitats: High versus low pesticide-input agroecosystems. Biological Control 152, <u>https://doi.org/10.1016/j.</u> <u>biocontrol.2020.104457</u>

Tillman, P. G. 2010. Parasitism and predation of stink bug (Heteroptera: Pentatomidae) eggs in Georgia corn fields. Environmental Entomology 39, <u>https://doi.org/10.1603/EN09323</u>.

Tillman, P. G., Toews, M., Blaauw, B., Sial, A., Cottrell, T., Talamas, E., *et al.* (2020) Parasitism and predation of sentinel eggs of the invasive brown marmorated stink bug, Halyomorpha halys (Stål) (Hemiptera: Pentatomidae), in the southeastern US. Biological Control, 145, <u>https://doi.org/10.1016/j.biocontrol.2020.104247</u>

Venette, R.C. & Hutchison, W.D. (2021) Invasive Insect Species: Global Challenges, Strategies & Opportunities. Frontiers in Insect Science 1, <u>https://doi.org/10.3389/</u> <u>finsc.2021.650520</u>

Brookdale Farm Supplies

Toro Tempus Ag Controller, a revolution in automation

The toro tempus ag controller allows for full farm automation. Tempus Ag uses a LoRA radio signal to create a bubble which allows for system automation. 1 base station produces a LoRa bubble of 5,200 feet in diameter. Multiple base stations can be added to cover large areas over one network for the entire application. The base stations can be operated on Wifi or with a 4G wireless signal. It can run irrigation cycles as well as collect environmental data, allowing growers to adjust their irrigation schedules as needed. Tempus Ag can report on a variety of sensors; temperature, pressure, soil moisture, humidity and more. Tempus works in both an outdoor field setting and an indoor high tunnel or greenhouse environment, Tempus can send alerts via text or email at thresholds determined by the user.



Trellis at Brookdale Fruit Farm, Hollis NH

🕨 Brookdale 🖉

Farm Supplies

UNA PASSIONE TUTTA ITALIANA

TORO

EMPUS-AG



Brookdale Farm Supplies has partnered with Valente for distribution in the United States. Valente's concrete posts are prestressed, reinforced posts that are trapezoidal shaped, with four smooth sides and no edges. This design prevents wear on hail netting and coverings. Valente trellis can be used in apple orchards, cherry orchards, vineyards and more. The system can be designed three different ways; standard trellis support, tall trellis support for future netting, or tall trellis with hail or over coverings included.

Orchard Tubing available in 17, 18 and 20 mm, Call for current pricing

NETAFIM

Rivulis

38 Broad Street Hollis, NH 03049 603-465-2240 www.brookdalefruitfarm.com



BlueLine[®]



STS-1113 FGN 1/4 Pg Ad DEC 2021 4.75"w x 7"h